

Please visit <https://www.evidentlybetter.org/bulletins/suicide-prevention/> to view our webpage featuring key links and emerging reports about suicide prevention.

Studies

[Similarities and differences in the prevalence and risk factors of suicidal behavior between caregivers and people with dementia: a systematic review](#)

Self-related and care receiver-related factors should be assessed among caregivers of people with dementia to evaluate the risk of suicidal behavior. In addition, we recommend evaluating suicidal risk in people with dementia in the early phase of dementia when cognitive decline is less severe. However, as the majority of the selected studies had unclear risk of bias, future studies with improved methodologies are warranted to confirm our study findings.

[Doctors working in paediatrics and anaesthetics at higher risk of suicidality, according to Australian survey](#)

Summarises a secondary cross-sectional analysis of survey data from Australian consultant doctors that explored associations between different medical specialities and the occurrence of common mental disorders and suicidal ideation.

[Effect of gender identity on the association between gender dysphoria and suicidality via appearance anxiety among transgender and gender-diverse young people: moderated mediation study](#)

Among TGD young people, gender dysphoria is significantly associated with suicidality via appearance anxiety, with gender identity moderating the mediating effects. Diverse treatments should consider the heterogeneity of TGD subgroups, with the aim of limiting the tendency of gender dysphoria to trigger appearance anxiety, thus further buffering against the risk of suicide.

[Risk factors for the need for advanced care among prescription and over-the-counter drug overdose patients](#)

Alcohol, paracetamol use and abnormal vital signs on scene might be risk factors for advanced care among prescription drugs or OTC drugs overdose patients, and the PAV score may predict the need for advanced care.

[Family intimacy and adaptability and non-suicidal self-injury: a mediation analysis](#)

Enhancing psychological resilience, controlling depressive symptoms, and reducing depression severity among adolescents by improving their family intimacy and adaptability are conducive to preventing and mitigating their NSSI behaviors.

[Family intimacy and adaptability and non-suicidal self-injury: a mediation analysis](#)

Enhancing psychological resilience, controlling depressive symptoms, and reducing depression severity among adolescents by improving their family intimacy and adaptability are conducive to preventing and mitigating their NSSI behaviors.



perfect care
Suicide prevention
crisis research risk self-harm
assessment resources
statistics information
study



NHS

Mersey Care
NHS Foundation Trust

Community and Mental Health Services

20 March 2024

Resource

[Family intimacy and adaptability and non-suicidal self-injury: a mediation analysis](#)

Enhancing psychological resilience, controlling depressive symptoms, and reducing depression severity among adolescents by improving their family intimacy and adaptability are conducive to preventing and mitigating their NSSI behaviors.

Webinar

[Suicide and Autism Awareness Training - launch webinar](#)

A new Autism and Suicide Awareness Training will be launched at a free online event, Tuesday 2 April, 12pm.

