

Please visit <https://www.evidentlybetter.org/bulletins/suicide-prevention/> to view our webpage featuring key links and emerging reports about suicide prevention.

Studies

[An Empirically Based, Recovery-Orientated Framework for Safety Planning Interventions for Suicide Prevention: An Integrative Literature Review and Critical Meta-Autoethnographic Synthesis](#)

The findings challenge dominant biomedical approaches, offering a more personalised, recovery-focused understanding of safety planning interventions.

[Does More Mean Better? Treatment Intensity and Predictors of Clinically Significant Change in Suicidal Ideation During Dialectical Behaviour Therapy](#)

The present study represents the first systematic examination of DBT treatment intensity in relation to suicidal ideation outcomes and the first evaluation of PiCD personality domains as moderators of change in suicidal ideation. Suicidal ideation decreased significantly across all three intensity levels over 12 weeks of treatment, with no significant differences in trajectories across treatment intensity levels. Although rates of CSC also did not differ significantly across treatment intensity levels, the relatively small number of participants achieving CSC limits conclusions regarding differences in clinically meaningful recovery across tracks.

[Unraveling Suicidal Ideation and Behaviors in a Global Cohort of Individuals at Clinical High-Risk for Psychosis: Multivariable Correlates and Subthreshold Psychotic Symptoms](#)

Overall, SIB in CHR-P individuals reflects a multidimensional clinical profile extending beyond subthreshold psychotic symptoms alone. Shared correlates related to affective symptoms and substance use, together with the evidence of a continuum between ideation and behaviors, underscore the need for tailored assessment and intervention strategies.

[Modulating factors for suicide reattempt among adolescents: A network analysis](#)

This study highlights the complexity of suicide reattempt risk factors in adolescents, emphasizing the importance of indirect interactions rather than isolated predictors. Incorporating standardized self-report tools may enhance clinical assessment and prevention strategies. Through network analysis, this research offers an innovative perspective to understand the multivariate nature of suicide risk in adolescent populations.

[Social disconnection integrates genetic and proteomic risks in suicidal ideation and depression](#)

This study presents one of the first comprehensive evaluations of how multidimensional social connection, genetic predisposition, and proteomic profiles are jointly associated with the vulnerability of SI and MDD in mid-to-late life. Leveraging a large population cohort, we find that multidimensional social connection shows strong associations with MDD and SI.

[Beyond psychiatric diagnosis: detection-gap perspective on suicide decedents in a population-based cohort](#)

Nearly half of suicide decedents had no recorded psychiatric diagnosis, and conventional psychosocial markers demonstrated limited discriminative value under current measurement conditions. These findings support a detection-gap framework and highlight the need for suicide prevention strategies extending beyond diagnosis-based approaches towards broader, population-level detection.

[‘It wasn’t just me’. Community voices of hope in suicide prevention: mixed-methods study of a community-based crisis service](#)

CBSC services would benefit from the expansion of NHS partnerships to extend regional reach, standardised monitoring and follow-up, and the strategic use of outcome data to strengthen advocacy and secure sustainable funding. Embedding this community-based model firmly within wider health systems has the potential to enhance its impact, ensure equitable access and inform broader systemic developments in suicide prevention.

[Invalidating environments and adolescent non-suicidal self-injury: longitudinal cross-lagged test of negative urgency and distress tolerance](#)

The findings highlight impulsive response under distress as a central mechanism linking environmental invalidation to adolescent NSSI and identify negative urgency as a key intervention target.

[A systematic review on prediction models for self-harm & suicide](#)

Is there a better way to predict suicide risk? Findings from a systematic review on prediction models for self-harm and suicide.

Call for research participants

[Changing the narrative on suicide and revisiting nurse regulation](#)

What we want to understand through Study 5 is much more about the relationship between having a complaint made about you and suicidality and distress.

Report

[The impact of patients suicide on the mental health of psychotherapists](#)

Interestingly, the preventability and predictability of patient suicides were assessed differently depending on whether the general views were solicited or if the question specifically referred to the “most distressing” suicide. Generally, a very small proportion considered suicides to be completely unavoidable or unpredictable. However, in both cases, about one-third of respondents believed that the suicide they had witnessed was entirely unpreventable or entirely unpredictable.

News

[Premier League clubs unite for Together Against Suicide](#)

Premier League clubs to offer tailored support on home matchdays, helping fans access trusted support in safe, non-judgemental environments.

[Campaign for Kien's Law](#)

Merseyside Police are backing the Kien's Law campaign, which calls for domestic abuse related suicides to be recognised in law. The campaign follows a six month Merseyside pilot that uncovered previously hidden domestic abuse concerns after suspected suicides, aiming to improve investigations, accountability, learning and prevention, while supporting victims and families.



Mersey Care
NHS Foundation Trust

15 September 2026

[Warrington MP calls for urgent action following closure of Papyrus](#)

A WARRINGTON MP has called for urgent action following the closure of a suicide prevention charity based in the town. It comes after the shock closure of Papyrus, a Warrington-based charity which sought to prevent young suicide and entered administration earlier this week. Its helpline, Hopeline247, ceased operating immediately – just days before World Suicide Prevention Day.

